

Quantitative measurements on the duplex stability of 2,6-diaminopurine and 5-chloro-uracil nucleotides using enzymatically synthesized oligomers

Jörg D. Hoheisel and Hans Lehrach

Imperial Cancer Research Fund, PO Box 123, Lincoln's Inn Fields, London WC2A 3PX, UK

Received 30 August 1990

2,6-Diaminopurine and 5-chloro-uracil 2'-deoxynucleoside 5'-triphosphates were synthesized from their 2'-deoxynucleosides. Using a method of creating oligonucleotides by enzymatic primer extension, dodecanucleotides representing an *XbaI/SalI* site and the complementary *SalI/XbaI* site were generated containing these base modifications. Their duplex stability was quantitatively compared by thin-layer chromatography to oligomers containing 2'-deoxyadenosine and 2'-deoxythymidine. The two unmodified oligomers already showed significant differences in dissociation temperature and binding equilibrium. Substitution with 5-chloro-2'-deoxyuridine did not affect the dissociation temperature of either oligomer, the 2,6-diaminopurine, however, led to an increase of 1.8°C or 1.5°C per modified base, respectively. While in the *XbaI/SalI* oligomer both base modifications changed the binding equilibrium, the 2,6-diaminopurine by a factor of 1.32, the 5-chloro-uracil by 0.65, no such effect was found with the complementary oligomer.

Thin-layer chromatography, 2,6-Diaminopurine, 5-Chloro-uracil, Thermodynamics

1. INTRODUCTION

Hybridization of short oligonucleotide probes has become a major tool in molecular biology, particularly more recently progressing towards the use of very short oligomers. Studier [1], for instance, proposed an improved sequencing strategy, based on enzymatic primer extension with dideoxynucleotides but using octanucleotide primers. Another new application is the physical mapping of large genomes by 'oligomer fingerprinting' whose practical feasibility has been demonstrated [2]. Also, large scale sequencing using hybridization of genomic DNA fragments to an array of oligomers (e.g. all 65 536 octamers) has been proposed [3,4]. All these methods, but also standard procedures like the polymerase chain reaction, for instance, would gain from an improvement in duplex stability of oligonucleotides. A way to achieve this is the substitution of natural nucleotides by base modifications, like pyrimidines with a halogen at the C⁵-position, which reportedly influences the base stacking [5], and 2,6-diaminopurine with a third hydrogen bond in its base pairing with thymine [6,7] which is known to thermally stabilize DNA-duplexes [8,9]. Since both 2,6-diaminopurine and 5-chloro-uracil 2'-deoxynucleosides are not available as phosphoramidites for a standard chemical synthesis of oligonucleotides, their 5'-

triphosphates were synthesized, being an ample source of nucleotide for the generation of different oligomers bearing modified (or unmodified) bases by means of a conventional fill-in reaction. Dodecamers representing the respective strand of an *XbaI/SalI* site were used to quantitatively determine how the base substitutions affect the dissociation temperature (T_d) and the binding equilibrium.

2. MATERIALS AND METHODS

2.1. Synthesis of nucleoside triphosphates

5-Chloro-2'-deoxyuridine (5-Cl-dUrd) and 2,6-diaminopurine 2'-deoxynucleoside (2-NH₂-dAdo) were from Sigma. Phosphorylation was carried out following a protocol modified from Goody and Isakov [10] and Ludwig [11]. 50 mg dry 2-NH₂-dAdo was taken up in 500 µl dry triethyl phosphate stirring under argon. 25 µl POCl₃ were added and the mixture was incubated at -20°C. In the meantime, 1 mmol pyrophosphoric acid was dissolved in 0.95 ml tri-*n*-butylamine and 2 ml methanol and dried in a rotary evaporator. Subsequently it was dried by evaporation twice from 5 ml pyridine, with 70 µl tri-*n*-butylamine also added before the second time. Finally, it was dissolved in 2 ml dry dimethyl formamide.

After 90 min at -20°C the phosphorylation mixture was evaporated to remove excess POCl₃ and the tri-*n*-butylammonium pyrophosphate in dimethyl formamide was added. Incubation was for 1.5 min at room temperature. The reaction was stopped by addition of 5 ml 0.2 M triethylammonium bicarbonate (pH 7.6) and kept on ice for 4 h. For 5-Cl-dUrd the conditions were identical, but 50 µl POCl₃ were added and the phosphorylation was carried out at room temperature for 4 h.

After the hydrolysis, the mixture was evaporated, the pH adjusted to 7.5, and extracted with 1 volume diethyl ether. Separation of the

Correspondence address: J.D. Hoheisel, Imperial Cancer Research Fund, PO Box 123, Lincoln's Inn Fields, London WC2A 3PX, UK

used to generate the oligonucleotides, reverse sequencing primer (New England Biolabs) was annealed to single stranded pJDH118, the primer's 3' end 85 bp distant from the DNA-terminus created by the *Hind*III-cut before the exonuclease III reaction (see section 2). Extension in the presence of 40 μ M and 180 μ M of the respective modified nucleotide resulted in blunt-ended molecules and showed no difference in comparison to dATP or dTTP (results not presented).

Measuring the melting behaviour of oligonucleotides by TLC perpendicular to a continuous temperature gradient proved to be very useful, because a whole temperature range is examined in a single experiment and different probes can be used simultaneously (e.g. Fig. 2). Probe concentrations similar to those in actual filter hybridizations can be used, enabling a direct application of the results to real hybridization experiments [13].

Surprisingly, the two unmodified oligomers (Fig. 1) were found significantly different in both T_d and binding equilibrium (Fig. 3). Although identical in base composition and very similar in sequence, with only the two blocks of hexanucleotides having swapped position, the difference in T_d was determined being 7.9°C [34.0°C ($\pm 0.35^\circ$ C) for oligomer I to 41.9°C ($\pm 1.2^\circ$ C) for oligomer II], and 65% ($\pm 0.6\%$) of oligomer II were bound below 10°C, that is by a factor of 1.2 more than the 54% ($\pm 1.5\%$) of oligomer I.

An autoradiogram is shown in Fig. 2 of an actual TLC experiment comparing the stability of oligomer I containing either dAdo or 2-NH₂-dAdo. Directly from the intensities of the radioactivity remaining at the origins, it is obvious that the substitution of 2-NH₂-dAdo for dAdo affected the oligomer's stability. The T_d increased by 5.5°C from 34.0°C ($\pm 0.35^\circ$ C) of the unmodified oligomer to 39.5°C ($\pm 0.7^\circ$ C) (Fig. 3A).




Fig. 3. Melting curves of oligomer I (A,C) and oligomer II (B,D). The two upper panels show the effect of substitution of dAdo (●, ◆) by 2-NH₂-dAdo (○, ◇), the lower panels the melting curves obtained with 5-Cl-dUrd (▽, □) instead of dThd (●, ◆). Each curve is the result of at least three entirely independent experiments, each point representing the average of 9 to 15 measurements; for each third point, the standard deviation is given.

Additionally, the binding equilibrium was improved over the whole temperature range. Below 10°C, for instance, 71.3% ($\pm 0.6\%$) of the modified oligomer was bound in comparison to only 54% ($\pm 1.5\%$) of the unmodified one. Using the oligomer II, the absolute difference in T_d was smaller with 4.6°C (41.9°C ($\pm 1.2^\circ\text{C}$) to 46.5°C ($\pm 0.6^\circ\text{C}$)). The amount of oligonucleotide bound below 10°C, however, did not change by substitution with 2-NH₂-dAdo (Fig. 3B).

5-Cl-dUrd (Fig. 3C) had no significant effect on the T_d of either oligomer. However, in oligomer I the substitution of 5-Cl-dUrd for dThd at positions 3 and 8 of the dodecamer reduced the amount of oligomer bound below 10°C by a factor of 0.65 to 35.1% ($\pm 0.8\%$) (the first nucleotide of the dodecamer could not be substituted, because it originated from the primer molecule used for its synthesis [12]). Again, there was no such effect in the complementary oligomer II, with the binding equilibrium being unchanged by the base modification (Fig. 3D).

4. DISCUSSION

The combination of the relatively simple phosphorylation of nucleosides to generate nucleotide triphosphates and the creation of oligomers by a conventional fill-in reaction provides a system to easily synthesize oligomers containing base modifications. Its use is even more versatile, because all components necessary – nucleotide triphosphates, template/primer oligonucleotides, uracil-DNA glycosylase – are stable if stored appropriately (uracil-DNA glycosylase is meanwhile commercially available from Boehringer-Mannheim). Here, the system was used to measure the effect of 2-NH₂-dAdo and 5-Cl-dUrd on the duplex stability of dodecamers.

Using 2-NH₂-dAdo, a considerable improvement in the duplex stability of short oligomers is achieved. Its incorporation should allow either more stringent conditions for primer annealing, thereby improving the specificity of the duplex formation and suppressing background problems, or the use of shorter oligomers. The lack of any significant difference in their effect on T_d between the halogen substitution of 5-Cl-dUrd and the 5-methyl group of dThd is in agreement with similar data obtained for the 5-bromo and 5-methyl derivatives of deoxycytidine [13].

The occurrence of changes in the binding equilibrium appears to be dependent on the sequence. This, and the considerable difference in duplex stability of the two unmodified dodecamers is the more surprising, since according to data from nearest neighbour analysis [16] the only difference in the sequence of the two oligomers (the central dinucleotide 'AG' or 'CT', respectively) should not matter. Further investigations into this direction are necessary to make the duplex formation of very short oligomers understood.

Acknowledgements: We wish to thank Roger S Goody and Marija Isakov for their help in establishing the technique of nucleoside phosphorylation in our laboratory, Iain Goldsmith for preparing the template/primer oligonucleotides, and Tomas Indahl for the uracil-DNA glycosylase. Funding by the Imperial Cancer Research Fund and a long-term EMBO fellowship to JH are gratefully acknowledged.

REFERENCES

- [1] Studier, F.W. (1989) *Proc. Natl. Acad. Sci. USA* 86, 6917–6921
- [2] Craig, A.G., Nizetic, D., Hoheisel, J.D., Zehetner, G. and Lehrach, H. (1990) *Nucleic Acids Res.* 18, 2653–2660.
- [3] Drmanac, R., Labat, I., Brukner, I. and Crkvenjakov, R. (1989) *Genomics* 4, 114–128.
- [4] Khrapko, K., Lysov, Y., Khorlyn, A., Shick, V., Florentiev, V. and Mirzabekov, A. (1989) *FEBS Lett.* 256, 118–122
- [5] Bugg, C.E., Thomas, J.M., Sundaralingam, M. and Rao, S.T. (1971) *Biopolymers* 10, 175–219.
- [6] Howard, F.B., Frazier, J. and Miles, H.T. (1976) *J. Biol. Chem.* 241, 4293–4295.
- [7] Rackwitz, H.R. and Scheit, K.H. (1977) *Eur. J. Biochem.* 72, 191–200.
- [8] Scheit, K.H. and Rackwitz, H.R. (1982) *Nucleic Acids Res.* 10, 4059–4069.
- [9] Chollet, A. and Kawashima, E. (1988) *Nucleic Acids Res.* 16, 305–317.
- [10] Goody, R.S. and Isakov, M. (1986) *Tetrahedron* 27, 3599–3602
- [11] Ludwig, J. (1987) in: *Biophosphates and Their Analogues × Synthesis, Structure, Metabolism and Activity* (Bruzik, K.S. and Stec, W.J. eds) pp. 201–204. Elsevier, Amsterdam, New York.
- [12] Craig, A.G., Nizetic, D. and Lehrach, H. (1989) *Nucleic Acids Res.* 17, 4605–4610.
- [13] Hoheisel, J.D., Craig, A.G. and Lehrach, H. (1990) *J. Biol. Chem.* 265, 16656–16660
- [14] Hoheisel, J.D. (1989) *Gene* 80, 151–154
- [15] Hoheisel, J.D. and Pohl, F.M. (1986) *Nucleic Acids Res.* 14, 3605.
- [16] Breslauer, K.J., Frank, R., Blocker, H. and Marky, L.A. (1986) *Proc. Natl. Acad. Sci. USA* 83, 3746–3750.